

RESULTS OBTAINED FROM SWEET CHERRY BREEDING IN IAȘI, ROMANIA

Elena IUREA¹, Sorina SÎRBU^{1,*},
Gelu CORNEANU¹, Margareta CORNEANU¹

*E-mail: sorinas66@yahoo.com

Received: Dec. 17, 2021. Revised: Feb. 21, 2022. Accepted: Mar. 09, 2022. Published online: Mar. 17, 2022

ABSTRACT. From 1994 - 2016, the Research Station for Fruit Growing (RSFG) Iași approved 28 cherry cultivars, of which three were early cultivars (Cetățuia, Cătălina and Amaris), 20 were middle maturation cultivars (Maria, Golia, Ștefan, Bucium, Iașrom, Tereza, Lucia, Radu, Oana, Alexus, Andreiaș, Ludovan, Cociuvaș, Iosifan, Paulică, Mihailis, Mușatini, Elaiși, Amar Maxut, and Amar Galata), and five had late maturation (Andante, Marina, Croma, Margonia, and George). Van, as the most widespread cultivar in Romanian orchards, was used as a control. The sweet cherry cultivars were created through direct hybridisation or free pollination. Maria Cultivar is the first Romanian self-fertile cultivar. The locally approved bitter cherry biotypes are Amar Maxut with black fruit, Amar Galata with double-coloured fruit, and Amaris with dark red fruit. Regarding the tree's vigour, cultivars Amaris, Tereza, Ștefan, and Golia recorded weak vigour, in comparison with the other cultivars, which had medium vigour. Bitter-tasting cherry cultivars have semi-firm pulp with semi-adherence to the stone, and the fruit weight was under 7 g. All 28 cultivars presented high precocity, productivity, fruit quality, and resistance to frost, drought, and diseases.

Keywords: cultivars; fruit; hybridisation; maturation stages; traits.

INTRODUCTION

In 1981, the Research Station for Fruit Growing (RSFG) Iași started collecting, preserving and studying cherry germplasm to use as initial material for cultivar registration. The Romanian bank of genes, represented by the national cherry collection, has 555 native and international genotypes, with old and new cultivars, local cherry biotypes, hybrids, and clones (Branîște *et al.*, 2007). The international objectives of the breeding programme were to improve the assortment of cherry cultivars with attractive qualities and genetic resistance to diseases, pests, and stressors (drought, frost) (Sansavini and Lugli, 2008), that reduce the vigour of trees and fruit of superior quality commercially, technologically, and in chemical traits, with ripening periods situated at the extremities of the cherry maturation season (Budan *et al.*, 1993; Budan and Grădinariu, 2000; Cociu,

¹ Research Station for Fruit Growing, Iași, Romania

1990; Quero-García *et al.*, 2017). In the last 20 years, the growers have established new and modern orchards with different cultivars with good fruits quality, low vigour for higher density of trees and suitable for climate conditions from the area (Sansavini and Lugli, 2008; Asănică *et al.*, 2013).

From 1994 - 2016, RSFG Iași registered 28 cherry cultivars in the catalogue of the State Institute for Testing and Registration of Cultivars, of which 19 represent 66% of the cultivars that get to remain in the assortment (Popescu *et al.*, 2015). Traditional methods were used to generate the new cultivars, namely hybridisation or free pollination, followed by harvesting of the hybrid stones, obtaining saplings, and selecting and testing the elite hybrids (Cociu and Oprea, 1989). It takes 15 - 23 years for a cultivar to go from hybridisation to registration due to the biological particularities of the species, including weak germination of the stones, weak viability of the saplings in the first years after being planted or transplanted, and late fruit entry. This paper presents the valuable features of the cherry cultivars created in RSFG Iași, that improve the native assortment and cultivars that have different ages of fruit maturation spread across the entire cherry maturation season.

MATERIALS AND METHODS

Soil and climate conditions

The natural environment where the research took place is located in the forest steppe of Moldova, where lands are slightly tilted from the northwest to the southeast, with an average slope of 5% and an altitude of 165 m. The soil is leached chernozem, weakly eroded by loess deposits and clay, with

a clay and sandy clay texture, a humus content between 2.5 and 3.5%, a pH of 5.8 - 6.8, a total average nitrogen content of 0.143 - 0.170, mobile phosphorus content of 20 - 82 ppm, and mobile potassium content of 188 - 360 K ppm. The annual average temperature (the average of the last 3 years) varies between 9.8 and 10.1°C. The highest monthly average temperature (21.5 - 21.8°C) is recorded in July, while January has the lowest temperature records. The average rainfall quantity recorded over a year varies between 520.8 and 603.5 mm, and the average atmospheric humidity is 70.2% (Corneanu *et al.*, 2007).

Plant material

The study took place between 2011 and 2016 and comprised 29 sweet cherry cultivars, of which 28 were approved at RSFG Iași (three bitter cherry cultivars and 25 sweet cherry cultivars), and Van was used as the control cultivar. The genotypes approved at RSFG Iași have the following genitors: Cetățuia (Van × Boambe de Cotnari); Cătălina (Van × Boambe de Cotnari); Alexis (Lijana OP); Andreiaș (HC 24/4 × Boambe de Cotnari); Ludovan (Van × Boambe de Cotnari), Lucia (Van × Muncheberge fruhe); Cociuș (Van × Bigarreau Moreau); Iosifan (Van × Lapins); Bucium (Van × Boambe de Cotnari); Maria (Van × Stella); Paulică (Bigarreau Drogan × Fromm); Ștefan (Van × Boambe de Cotnari); Tereza (Van × Ebony); Golia (Van × Boambe de Cotnari); Oana (Van × Boambe de Cotnari); Mihailis (Pietroase Geoagiu × New York 9801); Iașirom (Van × Boambe de Cotnari); Elaiși (Boambe de Cotnari × HC 21/1); Mușatini (Bigarreau Drogan OP); Radu (Van × Boambe de Cotnari); Andante (Lijana OP); Marina (Boambe de Cotnari × HC 23/31); Croma (Bigarreau Drogan × Van); Margonia (Van OP); and George (Ciliegia di Ottobre × Fromm). The capitalisation of the biological fund of existing genotypes in the spontaneous and cultivated flora in the Iași area was aided by

RESULTS OBTAINED FROM SWEET CHERRY BREEDING IN IAȘI, ROMANIA

positive selection of the valuable bitter cherry genotypes. Three genotypes of bitter cherry were selected and planted in the national collection and in the trial field 5×4 within RSFG Iași, two of which were approved as new varieties in 1994 under the names Amar Maxut and Amar Galata and one genotype that was approved in 2016 with the name Amaris. Bitter cherry genotypes have different ripening times: early (Amaris), middle (Amar Maxut), and late (Amar Galata). We evaluated nine trees (three trees in three repetitions) grafted on *Prunus mahaleb* L. in experimental plots and planted at m, with the crown guided as the free flattened palmette in the direction of the row with trees, without supporting and irrigation systems. On the row of trees, the soil was prepared with a rotary orchard tiller, and between the rows, the soil was grassed. Diseases and pests were controlled using phytosanitary treatments in accordance with the warnings received.

Determinations and analyses

To estimate resistance to anthracnosis (*Coccomyces hiemalis* Higg.), 300 leaves were observed, determining the frequency of the attack ($F\% = \text{number of attacked leaves from the total observed leaves}$), the intensity of the attack ($I\% = \text{percentage of attacked leaves of the total observed leaves}$: $I\% = (i \times f)/n$, where $i = \%$, and the leaf is attacked according to the mark in percentage; $f = \text{number of attacked leaves for each percentage from each mark}$, $n = \text{total number of attacked leaves}$), and the attack degree (GA), that represents the leaf attack, $GA\% = F \times I/100$ (Roșca *et al.*, 2011). To assess winter frost resistance, the viability of 100 flower buds from each third of the crown, for each variety, was analysed (Larsen, 2010). The phenophases of bloom and fructification were determined using BBCH monography as follows: 61 - beginning of flowering: about 10% of flowers open; 87 - fruit ripe for picking (Meier, 2001). The physical traits fruit and

stone weight (g) were measured by weighing 10 fruit/stones three times with the Radwag electronic scale with 0.01 g precision. The equatorial diameter of the fruit (mm) was determined using Luummytools digital calipers in 10 fruit three times. The soluble dry substance (SDS) was determined by measuring the refractive index with a Zeiss refractometer. Pulp firmness, fruit shape, and stone adherence to pulp were determined according to the UPOV TG/35/7 questionnaire for sweet cherry (UPOV, 2006). The fruit's resistance to cracking was determined using the Cristensen method, counting the number of cracked fruit after immersion in distilled water for six hours at 20°C (Webster and Looney, 1996).

The experimental data were statistically interpreted by XLSTAT 2021.5 software using the multiple comparisons method (Duncan test, with P 5%).

RESULTS AND DISCUSSION

Tree vigour, phenology and resistance of cherry cultivars

Cultivars with late fruit maturation were obtained using the following cultivars as genitors: Boambe de Cotnari (clones), Van, Fromm, Lijana, Bigarreau Drogan, and Ciliegia di Ottobre.

The breeding programme was continued to date, with the objectives being respected during development. Thus, cultivars with a ripening stage at the extremities of the cherry maturation season were obtained as early (Cetățuia, Cătălina, and Amaris) and late fruit maturation cultivars (Andante, Marina, Croma, Margonia, and George). The cherry cultivars are described in terms of vigour, bloom period, maturation age, qualitative traits of the fruit, and resistance to anthracnosis (*Coccomyces hiemalis* Higg.) and winter frost in

Tables 1 and 2. Regarding the tree's vigour, cultivars Amaris, Tereza, Ștefan, and Golia had weak vigour in comparison with the other cultivars, which had medium vigour.

Table 1 – Observations and measurements of 29 cherry cultivars (RSFG Iași, 2011 - 2016, n = 6)

Cultivar	Tree vigour ¹	Limit data (from earliest to latest/6 years):		Flower bud damage ³ (%)	Resistance to anthracnosis (<i>Coccomyces hiemalis</i> Higg.)		
		Start bloom (61; data) ²	Ripening time (87; data)		Frequency of attack (%)	Attack intensity (%) ⁴	Attack degree (%)
Cetățuia	5	03.04 - 20.04	11.05 - 30.05	6.0 ^{cde}	2.6	5	0.13
Amaris	3	06.04 - 19.04	23.05 - 04.06	9.0 ^c	3.0	4	0.12
Cătălina	5	05.04 - 21.04	17.05 - 04.06	8.0 ^{cd}	3.2	4	0.13
Alexus	5	14.04 - 24.04	10.06 - 23.06	28.0 ^a	3.7	4	0.15
Andreiaș	5	06.04 - 23.04	06.06 - 20.06	1.0 ^f	3.5	3	0.11
Ludovan	5	06.04 - 23.04	10.06 - 22.06	2.0 ^{ef}	2.9	4	0.12
Lucia	5	08.04 - 24.04	06.06 - 19.06	16.0 ^{bc}	3.1	4	0.12
Cociuvaș	5	05.04 - 23.04	09.06 - 20.06	2.0 ^{ef}	3.4	4	0.14
Iosifan	5	07.04 - 23.04	11.06 - 18.06	4.0 ^{def}	3.0	4	0.12
Bucium	5	06.04 - 23.04	07.06 - 18.06	6.0 ^{cde}	2.5	4	0.10
Maria	5	04.04 - 21.04	06.06 - 20.06	2.0 ^{ef}	3.1	4	0.12
Van	5	04.04 - 22.04	11.06 - 20.06	20.7 ^{ab}	2.5	5	0.13
Paulică	5	03.04 - 20.04	15.06 - 19.06	8.0 ^{cd}	2.9	3	0.10
Ștefan	3	07.04 - 25.04	06.06 - 17.06	2.0 ^{ef}	3.4	4	0.14
Tereza	3	05.04 - 24.04	09.06 - 17.06	2.0 ^{ef}	3.2	4	0.13
Golia	3	05.04 - 22.04	06.06 - 8.06	2.0 ^{ef}	3.9	4	0.16
Oana	5	02.04 - 21.04	05.06 - 19.06	4.0 ^{def}	3.7	4	0.15
Mihailis	5	05.04 - 23.04	05.06 - 16.06	2.0 ^{ef}	3.9	4	0.16
Iașirom	5	05.04 - 22.04	06.06 - 18.06	6.0 ^{cde}	2.9	4	0.12
Elaiși	5	05.04 - 22.04	13.06 - 20.06	0.0 ^f	3.6	4	0.14
Mușatini	5	08.04 - 23.04	08.06 - 20.06	2.0 ^{ef}	3.3	4	0.13
Amar Maxut	5	05.04 - 25.04	10.06 - 17.06	2.0 ^{ef}	2.9	4	0.12
Radu	5	02.04 - 20.04	05.06 - 16.06	8.0 ^{cd}	3.7	4	0.15
Amar Galata	5	05.04 - 25.04	10.06 - 23.06	1.0 ^f	2.7	4	0.11
Andante	5	07.04 - 24.04	18.06 - 06.07	4.0 ^{def}	3.4	4	0.14
Marina	5	04.04 - 23.04	17.06 - 06.07	24.0 ^{ab}	3.1	4	0.12
Croma	5	03.04 - 24.04	14.06 - 04.07	8.0 ^{cd}	3.0	4	0.12
Margonia	5	09.04 - 26.04	18.06 - 06.07	1.0 ^f	3.1	4	0.12
George	5	07.04 - 23.04	02.07 - 10.07	1.0 ^f	2.9	3	0.10

¹ UPOV test: tree's vigour mark on a scale of 1 - 9: 3 = weak; 5 = average (UPOV, 2006);

² BBCH phenological phases: 61-start bloom: about 10% of flowers open; 87-fruit ripe for picking (Meier, 2001);

³ Different letters correspond with a significant statistical difference at $P \leq 5\%$, Duncan test.

⁴ The attack intensity mark (1 - 6 scale): 1 = 1 - 3% attacked surface; 2 = 4 - 10%; 3 = 11 - 25%; 4 = 26 - 50%; 5 = 51 - 75%; 6 = 76 - 100% (Cociu and Oprea, 1989).

In regards to disease resistance, 2013 and 2016 were rainy years (with a surplus of 146.3 mm in 2013 and 128.2 mm in 2016), which is favourable for pathogen development; the cultivars expressed a low sensitivity to *Coccomyces hiemalis* Higg. The attack frequency was between 2.6 and 3.9% (*Table 1*).

Generally, in the six years of the study (2011 - 2016), during the dormancy period, the winters were mild, without extreme temperatures, but on February 12, 2012, - 24.3°C was recorded as the minimum temperature. The winter of 2013 was mild, but between March 15 and 29, the time when the cherry was out of dormancy, minimum temperatures were recorded at - 10.8°C (on March 24). Under these conditions, the flower buds are more sensitive, affecting the pistil inside the flower bud. The prolonged duration of these extremely low temperatures caused some varieties of cherries to suffer, causing some degree of damage to the flower buds, ranging from 0 to 28%, with the highest percentage in Alexis, Cetățuia, and Van (*Table 1*). However, flower bud damage up to 30% does not represent a crop danger (Wenden *et al.*, 2017). Resistance to frost is an important parameter for fruit tree species and varies from one cultivar to another, as well as within trees and the area of the tree (Szabó *et al.*, 1996; Szymajda *et al.*, 2013; Asănică *et al.*, 2014). During the six years of the study (2011 - 2016), the start of flowering occurred between April 2nd and 26th (*Table 1*). Harvest time was registered in the first half of May for the early varieties (Cetățuia, Cătălina and Amaris), in the first 20 days of June for

the mid-season varieties (Alexus, Andreiaș, Ludovan, Lucia, Cociuvaș, Iosifan, Bucium, Maria, Paulică, Ștefan, Tereza, Golia, Oana, Mihailis, Iașirom, Elaiși, Mușatini, Radu, Amar Maxut, and Amar Galata) and from the end of June to the beginning of July for the late varieties (Andante, Marina, Croma, Margonia and George) (*Table 1*). Similar studies showed that the phenological periods for the same cherry genotypes were variable depending on the climatic conditions of each year (Darbyshire *et al.*, 2012), but the ranking of cultivars remained the same. Throughout the study period, the earliest ripening time took place May 11th (Cetățuia) and the latest on July 10th (George), providing a period with fresh fruit between 40 and 61 days. Cultivar Maria is the first and only Romanian self-fertile cultivar. The quality of cherry fruit is generally determined by fruit size, skin colour, firmness of the pulp, stone size, and stone adherence to the pulp (Quero-Garcia *et al.*, 2017). The new cultivars have different fruit colours, including yellow, bicolour, shiny red, dark red, and black (*Fig. 1*).

Fruit characteristics

The fruit weight was between 3.6 (Amar Maxut) and 5.5 g (Amaris) for bitter taste genotypes, but for all sweet cherry genotypes, fruit weight ranged between 5.4 (Cetățuia) and 9.3 g (Andreiaș) (*Table 2*).

Compared with Van, as the control cultivar, which is widespread in Romanian orchards, Alexis, Andreiaș, Ludovan, Lucia, Iosifan, Elaiși and Mușatini had greater fruit weights (*Table 2*). The stone weight was not positively correlated with fruit weight in

Andante, which had the largest stones, but medium fruit with significant statistically differences, compared with

other genotypes, such as Cociuș or Iosifan, which have small stones but large fruit weight.

Table 2 - Physical and organoleptic traits in 29 cherry cultivars (RSFG Iași, average 2011 - 2016, n = 6)

Cultivar	Fruit weight ¹ (g)	Stone weight (g)	Fruit equatorial diameter (mm)	SDS (%) ²	Cracked fruit (%)	Pulp firmness ³	Pulp adherence to stone ⁴	Taste ⁵	Epidermis colour ⁶	Fruit shape ⁷
Early ripening cherry cultivars										
Cetățuia	5.4 ^{hi}	0.20 ^{he}	21.2 ^{ghi}	15.1 ^c	16.17 ^{bc}	SF	SA	S	DR	K
Amaris	5.5 ^h	0.26 ^{gh}	20.7 ^{hij}	19.0 ^a	1.22 ^h	SF	NA	B	DR	H
Cătălina	7.2 ^{fg}	0.30 ^{def}	23.8 ^{cdefg}	18.0 ^{ab}	9.00 ^{de}	SF	NA	S	SR	H
Mid-season ripening cherry cultivars										
Alexus	9.3 ^a	0.33 ^{bcd}	25.8 ^{abcd}	17.0 ^{bc}	7.33 ^{efg}	F	NA	S	SR	H
Andreaș	8.9 ^{abc}	0.33 ^{bcd}	24.9 ^{bcdef}	18.3 ^{ab}	8.67 ^{def}	F	NA	VS	SR	H
Ludovan	9.1 ^{ab}	0.33 ^{bcd}	26.5 ^{ab}	18.6 ^{ab}	2.83 ^{gh}	F	NA	VS	SR	K
Lucia	8.6 ^{abc}	0.33 ^{bcd}	25.4 ^{abcde}	19.0 ^{ab}	4.67 ^{efgh}	F	NA	S	SR	H
Cociuș	8.3 ^{abcde}	0.30 ^{def}	25.1 ^{bcdef}	18.5 ^{ab}	3.17 ^{gh}	F	NA	VS	SR	K
Iosifan	9.2 ^{ab}	0.32 ^{cde}	26.2 ^{abc}	17.2 ^{abc}	4.17 ^{efgh}	F	NA	VS	SR	K
Bucium	8.0 ^{cdefg}	0.30 ^{def}	25.0 ^{bcdef}	17.8 ^{ab}	18.50 ^{ab}	F	NA	S	SR	H
Maria	7.3 ^{efg}	0.28 ^{ef}	23.8 ^{cdefg}	18.5 ^{ab}	8.67 ^{def}	F	NA	S	SR	H
Van	7.3 ^{efg}	0.30 ^{cdef}	23.4 ^{defgh}	17.5 ^{abc}	43.50 ^a	F	NA	S	SR	K
Paulică	8.0 ^{cdefg}	0.32 ^{cde}	25.2 ^{bcdef}	17.1 ^{abc}	2.83 ^{gh}	F	NA	S	BR	K
Ștefan	8.0 ^{cdefg}	0.33 ^{bcd}	24.5 ^{cdef}	18.4 ^{ab}	15.50 ^{bc}	F	NA	S	SR	H
Tereza	8.2 ^{bcdef}	0.32 ^{cde}	27.8 ^a	17.6 ^{ab}	5.67 ^{efgh}	F	NA	S	SR	H
Golia	7.1 ^g	0.27 ^{fgh}	23.8 ^{cdefg}	17.1 ^{abc}	7.33 ^{efg}	F	NA	S	SR	H
Oana	7.0 ^g	0.33 ^{bcd}	23.3 ^{defgh}	17.0 ^{bc}	3.33 ^{gh}	F	NA	S	SR	K
Mihailis	7.1 ^g	0.34 ^{bc}	22.3 ^{fgh}	17.4 ^{abc}	3.00 ^{gh}	F	NA	S	DR	H
Iașirom	7.2 ^{fg}	0.26 ^{gh}	23.7 ^{cdefgh}	17.8 ^{ab}	1.17 ^h	F	NA	S	SR	H
Eliași	9.0 ^{abc}	0.33 ^{bcd}	25.5 ^{abcde}	18.3 ^{ab}	2.28 ^{gh}	F	NA	VB	SR	H
Mușatini	8.4 ^{abc}	0.29 ^{def}	24.4 ^{cdef}	17.7 ^{ab}	3.12 ^{gh}	F	NA	S	SR	K
Amar Maxut	4.1 ^j	0.28 ^{ef}	18.2 ^j	19.0 ^{ab}	1.45 ^h	SF	SA	VB	B	H
Radu	7.1 ^g	0.30 ^{def}	23.1 ^{defgh}	17.0 ^{bc}	2.17 ^{gh}	F	NA	S	SR	K
Late ripening cherry cultivars										
Amar Galata	4.3 ^{ij}	0.34 ^{bc}	18.6 ^{ij}	18.4 ^{ab}	5.97 ^{efgh}	SF	SA	SB	BR	SE
Andante	8.3 ^{abcde}	0.38 ^a	25.7 ^{abcde}	18.5 ^{ab}	9.33 ^{de}	F	NA	S	Y	K
Marina	7.5 ^{defg}	0.33 ^{bcd}	24.1 ^{cdefg}	17.6 ^{ab}	12.83 ^{cd}	F	NA	S	BR	H
Croma	8.2 ^{bcdef}	0.33 ^{bcd}	24.7 ^{cdef}	18.3 ^{ab}	18.33 ^b	F	NA	S	SR	K
Margonia	7.1 ^g	0.36 ^{ab}	23.0 ^{defgh}	17.0 ^{bc}	1.88 ^{gh}	F	NA	S	Y	H
George	7.0 ^g	0.33 ^{bcd}	22.9 ^{efgh}	17.3 ^{abc}	3.00 ^{gh}	F	NA	S	SR	H

¹ Different letters correspond with the significant statistical difference for P ≤ 5%, Duncan test;

² soluble dry solids;

³ pulp firmness: F = firm; SF = semi-firm;

⁴ NA = non-adherent; SA = semi-adherent;

⁵ S = sweet; VS = very sweet; B = bitter; SB = slightly bitter;

⁶ DR = dark red; SR = shiny red; BR = bicoloured; B = black; Y = yellow;

⁷ H = heart-shaped; K = kidney-shaped; SE = spherical elongated.

RESULTS OBTAINED FROM SWEET CHERRY BREEDING IN IAȘI, ROMANIA



Figure 1 – Cherry cultivars obtained in RSFG Iași, with different maturation seasons, shapes and fruit skin colours:

1) Cetățuia; 2) Amar Galata; 3) Cătălina; 4) George; 5) Marina; 6) Andante (original)

For equatorial fruit diameter, the studied cultivars ranged between 18.2 (Amar Maxut) and 27.8 mm (Tereza). Additionally, the Ludovan cultivar had great fruit size with no significant statistical differences from Tereza. Cetățuia, Amaris, Amar Galata, and Amar Maxut had a small fruit size, but was considered a large fruit (4.1 - 5.5 g) for the early and bitter varieties. Budan (2014) showed that bitter and early varieties are smaller in size (weight and equatorial diameter) than medium or late ripening sweet cherries.

The soluble dry substance was between 15.1 (Cătălina) and 19.0% (Amaris, Ludovan, Amar Maxut). These results are consistent with other studies of cherry cultivars, in which significant variations were found among cultivars (Skrzyński *et al.*, 2016). The percentage of fruit cracking ranged from 1.17 (Iașirom) to 43.5% (Van). Compared with Van, all other studied cultivars

were more resistant to rain-induced fruit cracking (*Table 2*), which is a major physiological problem in sweet cherry cultivars (Michailidis *et al.*, 2020). All varieties had a firm pulp and were not adherent to the stone, except for the early and bitter varieties (Cetățuia, Cătălina, Amaris, Amar Maxut, and Amar Galata), in which the firmness of the pulp and adherence to the stone were average (*Table 2*).

Three varieties were bitter (Amaris, Amar Galata, Amar Maxut), and the remaining 25 cultivars were sweet. The bitter taste of fruit is an important parameter for producers of traditional products, such as liqueurs or jams (Budan, 2014). Regarding the shape of the fruit, 10 varieties had a kidney shape, namely Cetățuia, Ludovan, Cociuvaș, Iosifan, Paulică, Oana, Mușatini, Radu, Andante, Croma, and Van. One variety was oblate (Amar Galata), and all other varieties had a heart shape (*Table 2*).

CONCLUSIONS

The new cultivars approved at RSFG Iași can be recommended for early ripening time (Cetățuia and Cătălina), late flowering (Margonia), special fruit quality and late ripening time (Marina, Andante, Margonia, Croma, and George), and weak tree vigour (Golia, Tereza, and Ștefan). Andreiaș, Amar Galata, and George had good resistance to anthracnose and winter frost.

Alexus, Iosifan, Ludovan, Elaiăși, Andreiaș, Lucia, Mușatini, Cociuvaș, and Andante have large fruits and high soluble dry solids content, so they are recommended for fresh consumption or processing. The new cherry varieties introduced in the current assortment can guarantee the success of orchards in northeast Romania.

Acknowledgement. This work was partially supported by the Romanian Ministry of Agriculture and Development from Financial Programs ADER 7.2.2. with the title: "Implementation of new breeding methods of the fruit tree species to improve and reduce the time in the selection process".

REFERENCES

- Asănică, A., Tudor, V. & Teodorescu, R. (2013). Distinctive behaviour of some sweet cherry cultivars related to rootstock type. *AgroLife Sci. J.*, 2(1): 79-82.
- Asănică, A., Tudor, V. & Țiu, J.V. (2014). Frost resistance of some sweet cherry cultivars in the Bucharest area. *Sci. Papers. Ser. B, Hort.*, Vol. LVIII: 19-24.
- Asănică, A., Tudor, V. & Teodorescu, R. (2013). Distinctive behaviour of some sweet cherry cultivars related to rootstock type. *AgroLife Sci. J.*, 2(1): 79-82.
- Branîște, N., Budan, S., Butac, M. & Militaru, M. (2007). Varieties of trees, fruit bushes and strawberries created in Romania (in Romanian). *Paralela 45 Publishing House*, Pitești, 476 p.
- Budan, S. (2014). Traditional and commercial uses of Romanian bitter cherry cultivars. *Acta Hort.*, 1032: 25-28, DOI: 10.17660/ActaHortic.2014.1032.2
- Budan, S. & Grădinaru, G. (2000). The cherry (in Romanian). *Ion Ionescu de la Brad Publishing House*, Iași, 262 p.
- Budan, S., Iezzoni, A. & Andersen, R.L. (1993). Evaluation of some New York sweet cherry selections in Romania. *Fruit Var. J. (APS)*, 47: 168-171.
- Cociu, V. & Oprea, Ș. (1989). Research methods in fruit plant improvement (in Romanian). *Dacia Publishing House*, Cluj-Napoca, 172 p.
- Cociu, V. (1990). New varieties a factor in the progress of fruit growing. *Ceres Publishing House*, Bucharest, 323 p.
- Corneanu, G., Cârdei, E., Petre, L., Corneanu, M., Maxim, O. & Palade, I. (2007). Iași Fruit Research and Development Station - 30th Anniversary of Scientific Research and Development 1977-2007. *Altfel Publishing House*, 233 p.
- Darbyshire, R., Webb, L., Goodwin, I. & Barlow, E.W.R. (2012). Evaluation of recent trends in Australian pome fruit spring phenology. *Int. J. Biometeorol.*, 57(3): 409-421, DOI:10.1007/s00484-012-0567-1.
- Ganji Moghaddam, E., Ahmadi Moghaddam, H. & Piri, S. (2012). Genetic variation of selected Siah Mashhad sweet cherry genotypes grown under Mashhad environmental conditions in Iran. *Crop Breed. J.*, 3(1): 45-51, DOI: 10.22092/cbj.2013.100449.
- Głowacka, A. & Rozpara, E. (2014). Examination of the suitability of different pollinators for four sweet cherry cultivars commonly grown in Poland. *J. Hort. Res.*, 22(1): 85-91, DOI: 10.2478/johr-2014-0010.

- Larsen, H.J. (2010).** Evaluating tree fruit bud and fruit damage from cold. *Colorado State University Extension*, Fact Sheet No. 7.426, Gardening Series: Trees and Shrubs.
- Meier, U. (Ed.) (2001).** Growth stages of mono- and dicotyledonous plants - BBCH Monograph. *Julius Kühn-Institute (JKI)*, <http://www.jki.bund.de>
- Michailidis, M., Karagiannis, E., Tanou, G., Sarrou, E., Karamanoli, K., Lazaridou, A., Martens, S. & Molassiotis, A. (2020).** Sweet cherry fruit cracking: follow-up testing methods and cultivar-metabolic screening. *Plant Methods*, 16(51), DOI: 10.1186/s13007-020-00593-6
- Popescu, M., Ciora, M., Bran, A. & Theodorescu, M. (2015).** The official catalog of cultivated plant varieties in Romania for 2015, *MADR-ISTIS* Bucharest, 112 p.
- Quero-Garcia, J., Iezzoni, A., Pulawska, J. & Lang, G. (Eds.) (2017).** Cherries: botany, production and uses, *CABI*, Boston, MA, 550 p., ISBN: 9781780648378, DOI: 10.1079/9781780648378.0000
- Roșca, I., Oltea, I. & Mitrea, I. (2011).** General and special entomology treatise (in Romanian), *ALPHA MDN Publishing House*, Buzău, 1055 p.
- Sansavini, S. & Lugli, S. (2008).** Sweet cherry breeding programs in Europe and Asia. *Acta Hort.*, 795: 41-58, DOI: 10.17660/ActaHortic.2008.795.1
- Skrzyński, J., Leja, M., Gonkiewicz, A. & Banach, P. (2016).** Cultivar effect on the sweet cherry antioxidant and some chemical attributes. *Folia Hort.*, 28(1): 95-102, DOI: 10.1515/fhort-2016-0011
- Szabó, Z., Nyéki, J. & Soltész, M. (1996).** Frost injury to flower buds and flowers of cherry varieties. *Acta Hort.*, 410: 315-322, DOI: 10.17660/ActaHortic.1996.410.49
- Szymajda, M., Pruski, K., Żurawicz, E. & Sitarek, M. (2013).** Freezing injuries to flower buds and their influence on yield of apricot (*Prunus armeniaca* L.) and peach (*Prunus persica* L.). *Can. J. Plant Sci.*, 93(2): 191-198, DOI: 10.4141/cjps2012-238
- UPOV (2006).** Guidelines for the conduct of tests for distinctness, uniformity and stability. TG/35/7 - Sweet cherry (*Prunus avium* L.). *International Union for the Protection of New Cultivars of Plants*, Geneva, Switzerland, 31 p.
- Webster, A.D. & Looney, N.E. (Eds.) (1996).** Cherries: Crop Physiology, Production and Uses. *CABI*, Wallingford, UK, 513 p.
- Wenden, B., Campoy, J.A., Jensen, M. & Lopez-Ortega, G. (2017).** Climatic limiting factors: temperature. In: Cherries: Botany, Production and Use. J. Quero-Garcia, Iezzoni, A., Pulawska, J., Lang, G. (Eds.). *CABI*, Boston, MA., 166-189.